



11 Rec'd PCT 14 APR 1997

08/765,695
154-130
PCT
1/2/97

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:	§	
	§	
LARS ABRAHMSEN	§	
PER BJORK	§	Attorney Docket No.: A96335US
MIKAEL DOHLSTEN	§	
TERJE KALLAND	§	
	§	
	§	
Serial No. 08/765,695	§	Group Art Unit:
	§	
Filed: 10 January 1997	§	Examiner:
	§	
For: A CONJUGATE BETWEEN	§	
A MODIFIED SUPERANTIGEN	§	
AND A TARGET-SEEKING	§	
COMPOUND AND THE USE OF	§	
THE CONJUGATE	§	

**RESPONSE TO NOTICE TO FILE MISSING
PARTS OF APPLICATION UNDER 37 CFR 1.53(d)**

Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

In response to the notice mailed from the U.S. Patent Office on March 10, 1997 in
the above-referenced application, enclosed are the following:


1. Declaration of Lars Abrahmsen;
2. Declaration of Per Bjork and Mikael Dohlsten;
3. Declaration of Terje Kalland;
3. \$130.00 Requisite Fee for Filing Missing Parts; and
4. Copy of Form PCT/DO/EO/905.

520 KD 03/29/97 08765695
1 154 130.00 CR ✓

The Commissioner is respectfully petitioned for any extension of time necessary in order to render this response timely.

Please charge any fees due or credit any overpayment to Deposit Account Number 16-2435.

Respectfully submitted,



David L. Fox, Ph.D.

Registration No. P40,612

Date: April 9, 1997

PRAVEL, HEWITT, KIMBALL & KRIEGER
1177 West Loop South
Tenth Floor
Houston, TX 77027
(504)835-2000

CERTIFICATION UNDER 37 CFR 1.8(a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, on April 9, 1997.



David L. Fox, Ph.D.

Registration No. P40,612

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

A96335US

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

08/765695

INTERNATIONAL APPLICATION NO

PCT/SE95/00681

INTERNATIONAL FILING DATE

7 June 1995 (07.06.95)

PRIORITY DATE CLAIMED

11 July 1994 (11.07.94)

TITLE OF INVENTION A CONJUGATE BETWEEN A MODIFIED SUPERANTIGEN AND A TARGET-SEEKING
COMPOUND AND THE USE OF A CONJUGATE**APPLICANT(S) FOR DO/EO/US**

ABRAHMSEN, Lars; BJORK, Per; DOHLSTEN, Mikael; KALLAND, Terje

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendemnts has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

FEE VALUE
FOCUS NUMBER
DEPOSIT ACCOUNT NO.
76-2733
DATE RECEIVED
08-04-95
FILED
U.S. PATENT & TRADEMARK OFFICE
PHOENIX, ARIZONA

17. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):**

Search Report has been prepared by the EPO or JPO. \$880.00

International preliminary examination fee paid to USPTO (37 CFR 1.482) \$680.00

No international preliminary examination fee paid to USPTO (37 CFR 1.482)
but international search fee paid to USPTO (37 CFR 1.445(a)(2)) \$750.00Neither international preliminary examination fee (37 CFR 1.482) nor
international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$1010.00International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(2)-(4) \$94.00**ENTER APPROPRIATE BASIC FEE AMOUNT =**

\$ 1,010

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	38 - 20 =	18	X \$22.00
Independent claims	2 - 3 =	0	X \$78.00

\$ 396

\$ -0-

MULTIPLE DEPENDENT CLAIM(S) (if applicable)

+ \$250.00

\$

TOTAL OF ABOVE CALCULATIONS =

\$ 1,406

Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

\$

SUBTOTAL =

\$

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

TOTAL NATIONAL FEE =

\$

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

+

\$

TOTAL FEES ENCLOSED =

\$ 1,406

Amount to be:
refunded

\$

charged

\$

a. ☒ A check in the amount of \$ 1,406.00 to cover the above fees is enclosed.b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 16-2435. A duplicate copy of this sheet is enclosed.**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO

Paul E. Krieger
Pravel, Hewitt, Kimball & Krieger, P.C.
1177 West Loop South, Tenth Floor
Houston, Texas 77207-9095

SIGNATURE

Jan K. Simpson

NAME

33,283

REGISTRATION NUMBER

parent PCT application. The declarations submitted herewith are facsimiled copies. Originals of the declarations will be sent to the Patent & Trademark Office as soon as they are received from Sweden.

Amendment

IN THE CLAIMS:

Please cancel claims 1-13 without prejudice and please insert in their place the following substitute claims that have been numbered beginning with number one.

Therefore, please add the following new claims:

1. A conjugate comprising:
 - a. a biospecific affinity counterpart that is capable of binding to a predetermined structure, and
 - b. a peptide that
 - i. contains an amino acid sequence that is derived from a superantigen,
 - ii. has the ability to bind to a V β of a T cell receptor, and
 - iii. has been mutated to show a modified ability to bind to MHC class II antigens compared to the superantigen from which the peptide is derived,

which parts are covalently linked together.

2. The conjugate according to claim 1 wherein:
 - a. the biospecific affinity counterpart is directed towards a cell surface structure, and
 - b. the conjugate has the ability to activate T-lymphocytes to lyse cells that exhibit the cell surface structure on their cell surface.
3. The conjugate according to claim 2, wherein the biospecific counterpart is directed against a cell surface structure associated with a disease.
4. The conjugate according to claim 3, wherein the disease is selected from the group consisting of cancers, viral infections, autoimmune diseases and parasitic infestations.
5. The conjugate according to claim 4, wherein the disease is a cancer.
6. The conjugate according to claim 5, wherein the biospecific affinity counterpart comprises a polypeptide structure.
7. The conjugate according to claim 6, wherein the biospecific counterpart is selected from the group consisting of an antibody or an antigen-binding fragment thereof.

8. The conjugate according to claim 6, wherein the biospecific affinity counterpart and the peptide are fused together.
9. The conjugate according to claim 8, wherein the biospecific counterpart is selected from the group consisting of an antibody or an antigen-binding fragment thereof.
10. The conjugate according to claim 2, wherein the ability of binding to MHC class II has been altered by at least 10%.
11. The conjugate according to claim 6, wherein the ability of binding to MHC class II has been reduced.
12. The conjugate according to claim 9, wherein the ability of binding to MHC class II has been reduced.
13. The conjugate according to claim 2, wherein the superantigen is a staphylococcal enterotoxin.
14. The conjugate according to claim 6, wherein the superantigen is a staphylococcal enterotoxin.

15. The conjugate according to claim 9, wherein the superantigen is a staphylococcal enterotoxin.

16. The conjugate according to claim 10, wherein the superantigen is a staphylococcal enterotoxin.

17. The conjugate according to claim 11, wherein the superantigen is a staphylococcal enterotoxin.

18. The conjugate according to claim 12, wherein the superantigen is a staphylococcal enterotoxin.

19. The conjugate according to claim 2, wherein the superantigen is a superantigen requiring zinc ions for binding to MHC class II antigens.

20. The conjugate according to claim 19, wherein the superantigen has been mutated in a codon encoding an amino acid residue which coordinates zinc when the superantigen binds to MHC class II antigens.

21. The conjugate according to claim 20, wherein the superantigen is a staphylococcal enterotoxin.

22. The conjugate according to claim 21, wherein the superantigen is selected from the group consisting of staphylococcal enterotoxin A or E.

23. A method for the treatment of a diseased condition in a mammal, which condition means the presence of specific cells that are associated with the condition by the expression of a disease specific cell surface structure, wherein one administers to the mammal a therapeutically effective amount of covalent conjugate that is able to activate T lymphocytes to lyse cells that carry the disease specific cell surface structure and comprises:

- a. a biospecific affinity counterpart that is capable of binding to said surface structure, and
- b. a peptide that
 - i. contains an amino acid sequence that is derived from a superantigen,
 - ii. has the ability to bind to a $V\beta$ of a T cell receptor, and
 - iii. has been mutated to show a modified ability to bind to MHC class II antigens compared to the superantigen from which the peptide is derived.

24. The method of claim 23, wherein the disease is selected from the group consisting of cancers, viral infections, autoimmune diseases and parasitic infestations.

25. The method of claim 24, wherein the disease is a cancer.
26. The method of claim 23, wherein the superantigen is a superantigen requiring zinc ions for binding to MHC class II antigens.
27. The method of claim 26, wherein the superantigen has been mutated in a codon encoding an amino acid residue which coordinates zinc when the superantigen binds to MHC class II antigens.
28. The method of claim 23, wherein the superantigen is a staphylococcal enterotoxin.
29. The conjugate according to claim 23, wherein the superantigen is selected from the group consisting of staphylococcal enterotoxin A or E.
30. The method of claim 27, wherein the superantigen is a staphylococcal enterotoxin.
31. The method of claim 23, wherein the biospecific affinity counterpart comprises polypeptide structure.

32. The method of claim 31, wherein the biospecific affinity counterpart is selected from the group consisting of an antibody or an antigen-binding fragment thereof.

33. The method of claim 31, wherein the biospecific counterpart and the peptide are fused together.

34. The method of claim 32, wherein the biospecific counterpart and the peptide are fused together.

35. The method of claim 34, wherein the superantigen is a superantigen requiring zinc ions for binding to MHC class II antigens.

36. The method claim 35, wherein the superantigen has been mutated in a codon encoding an amino acid residue which coordinates zinc when the superantigen binds to MHC class II antigens.


37. The method of claim 35, wherein the superantigen is a staphylococcal enterotoxin.

38. The conjugate according to claim 37, wherein the superantigen is selected from the group consisting of staphylococcal enterotoxin A or E.

REMARKS

Please add the above-listed new and substitute claims to the National Phase Application of the above-referenced PCT application. Please begin claim numbering with the number one. The above-noted new and substitute claims reflect amendments to the claims of the parent PCT application and do not include new matter.

Respectfully submitted,


Jan K. Simpson
Reg. No. 33,283

PRAVEL, HEWITT, KIMBALL & KRIEGER
1177 West Loop South
Tenth Floor
Houston, TX 77027
(713) 850-0909

Date: 1/10/97

18 hcd PCT/PTO 10 JAN 1997

1

A CONJUGATE BETWEEN A MODIFIED SUPERANTIGEN AND A TARGET-SEEKING
COMPOUND AND THE USE OF THE CONJUGATE.

Superantigens are primarily proteins of viral or bacterial
5 origin and are capable of simultaneous binding to MHC class II
antigens on mammalian cells and the T cell receptor V β chain. The
binding leads to activation of T-lymphocytes and lysis of the MHC
class II bearing cells. The moderate degree of polymorphism of
the binding part of the V β chain causes a relatively large
10 portion of the T-lymphocytes to be activated when contacted with
a superantigen (in comparison with activation through normal
antigen-processing).

Initially the superantigen concept was associated with various
staphylococcal enterotoxins (SEA, SEB, SEC₁, SEC₂, SED, and SEE).
15 Recently a new staphylococcal enterotoxin named SEH has been
discovered (Keyong et al., J. Exp. Med. 180 (1994) 1675-1683).
After the interest had been raised, further superantigens were
discovered. Examples are Toxic Shock Syndrome Toxin 1 (TSST-1),
Exfoliating Toxins (Exft) that are associated with scalded skin
20 syndrome, Streptococcal Pyrogenic Exotoxin A, B and C (SPE A, B,
and C), Mouse Mammary Tumor Virus Proteins (MMTV), Streptococcal
M Proteins, Clostridial perfringens enterotoxin (CPET) among
others. For a review of superantigens and their properties see
Kotzin et al. (Adv. Immunol. 54 (1993) 99-166).

25 Pseudomonas exotoxin A has been looked upon as a functional
superantigen because there are results indicating that this toxin
may be processed intracellularly by accessory cells to fragments
that are expressed on the cell surface with the ability to bind
to the V β chain and a subsequent activation of T cells.
30 (Pseudomonas exotoxin A. Legaard et al., Cell. Immunol. 135
(1991) 372-382).

Superantigens as such have been suggested for therapy of
various diseases with curative effects being accomplished through
a general activation of the immune system (Kalland et al., WO
35 9104053; Terman et al., WO 9110680; Terman et al., WO 9324136;
Newell et al., Proc. Natl. Acad. Sci. USA 88 (1991) 1074-1078).

CONFIRMATION
COPY

In connection with vaccines it has been suggested to use superantigens that have been mutated so as to lose their TCR binding ability (Kappler & Marrack, WO 9314634).

The mutation of superantigens has previously been described
5 (Kappler & Marrack, WO 9314634; Kappler et al., J. Exp. Med. 175 (1992) 387-396; Grossman et al., J. Immunol. 147 (1991) 3274-3281; Hufnagle et al., Infect. Immun. 59 (1991) 2126-2134).

We ourselves have previously suggested to employ conjugates between a superantigen and an antibody for therapy in order to
10 lyse cells that express the structure towards which the antibody is directed (Dohlsten et al., WO 9201470; Lando et al., Cancer Immunol. Immunother. 36 (1993) 223-228; Kalland et al., Med. Oncol. Tumor Pharmacother. 10 (1993) 37-47; Lando et al., J. Immunol. 150 (8 part 2) (1993) 114A (Joint Meeting of the
15 American Association of Immunologists and the Clinical Immunology Society, Denver, Colorado, USA, May 21-25 (1993)); Lando et al., Proc. Am. Assoc. Cancer Res. Annu. Meet. 33(0) (1992) 339 (Annual meeting of the American Association for Cancer Research, San Diego, California, USA, May 20-23 (1992)); Dohlsten et al., Proc.
20 Natl. Acad. Sci. USA 88 (1991) 9287-9291). Diseases suggested to be treated have been cancers, viral infections, parasitic infestations, autoimmune diseases and other diseases associated with cells expressing disease-specific surface structures. The experimental work carried out so far has focused on conjugates
25 containing recombinant SEA and various anti-cancer antibodies. The conjugates as such have had a somewhat reduced ability to bind MHC class II antigens compared to the non-conjugated form of the superantigen. It has not been determined if a decreased MHC class II antigen binding ability is beneficial or not for
30 achieving an optimal lyse and an optimal therapeutic effect.

Immune therapy experiments with SEB chemically conjugated to a tumor specific anti-idiotypic antibody have previously been described by Ochi et al., (J. Immunol. 151 (1993) 3180-3186).

During the prosecution of the priority application the Swedish
35 Patent Office has additionally cited Buelow et al. (J. Immunol. 148 (1992) 1-6) that describes fusions between Protein A and

fragments of SEB without emphasis of the MHC classes II binding or use of the fusion for cell killing; and Hartwig et al. (Int. Immunol. 5 (1993) 869-875) that describes mutations affecting MHC class II binding of the non-fused form of the superantigen streptococcal erythrogenic toxin A.

The objectives of the invention

A first objective of the invention is to improve previously known superantigen-antibody conjugates with respect to general immune stimulation versus directed cytotoxicity. Stimulation results in activated T-lymphocytes and is dependent on the ability of the superantigen to bind to both the T cell receptor and an MHC class II antigen.

A second objective of the invention is to provide conjugates between biospecific affinity counterparts (e.g. antibodies) and superantigens with a modified affinity for MHC class II antigens. This has now been shown to improve the selectivity for superantigen antibody dependent cell cytolysis (SADCC) of cells exposing the antigen (against which the antibody/biospecific affinity counterpart of the conjugate is directed) over other cells exposing MHC class II antigens.

A third objective of the invention is to provide conjugates that can be used as the active principle in the treatment of mammals suffering from cancers, autoimmune diseases, parasitic infestations, viral infections or other diseases associated with cells that on their surface express structures that are specific for respective disease.

The invention

- The main aspect of the invention is a conjugate comprising
- a. a biospecific affinity counterpart that is directed towards a structure to which one intends to bind to the conjugate,
 - b. a peptide that
 - i. is derived from a superantigen,
 - ii. has the ability to bind to the V β chain of the T cell receptor, and

iii. has a modified ability to bind to MHC class II antigens compared to the superantigen from which the peptide is derived (wild-type of superantigen = SA(wt)).

5 The peptide and the affinity counterpart are covalently linked to each other via a bridge (B).

The preferred conjugates have the ability to activate and direct T-lymphocytes to selective lysis of cells that on their surface expose the structure against which the affinity
10 counterpart is directed. This means that the conjugates shall cause cytolysis in an SADCC mediated method (Superantigen Antibody Dependent Cellular Cytotoxicity). See the experimental part below and our previous publications concerning conjugates between superantigens and antibodies (e.g. Dohlsten et al., WO
15 9201470).

The inventive conjugates have a structure that is analogous to the superantigen-antibody conjugates described in the prior art (Dohlsten et al., WO 9201470 which hereby is incorporated by reference), i.e. the conjugates complies with the formula:

20
$$T-B-SA(m)$$

where T represents the biospecific affinity counterpart, SA(m) is the modified superantigen (the above-mentioned peptide), and B is a covalent bridge linking T and SA(m) together.

T can in principle be any structure that binds via biospecific
25 affinity. In most important cases, T is capable of binding to a cell surface structure, preferably a disease specific structure as given above. The structure against which T is directed is usually different from (a) the V β chain epitope to which the superantigen derived peptide (SA(m)) binds and (b) the MHC class
30 II antigen epitope to which the unmodified superantigen binds. The biospecific affinity counterpart T may primarily be selected among interleukins (e.g. interleukin-2), hormones, antibodies and antigen binding fragments of antibodies, growth factors etc. See for instance Woodworth, Preclinical and Clinical Development of
35 Cytokine Toxins presented at the conference "Molecular Approaches to cancer Immunotherapy", Ashville, North Carolina, November 7-

11, 1993. Polypeptides binding to the constant domains of immunoglobulins (e.g. Proteins A and G and L), lectins, streptavidin, biotin etc were at the priority date considered to be of minor importance.

5 At the priority date, it was preferred that T was an antibody or an antigen binding fragment of an antibody (including Fab, F(ab)₂, Fv, single chain antibody etc), with particular emphasis of an antibody active fragment (such as Fab) of antibodies directed against the so called C242 epitope (Lindholm et al., WO 10 9301303) or against other cancer specific epitopes.

In case T is an antibody it is primarily monoclonal or a mixture of a defined number of monoclonals (e.g. 2, 3, 4, 5 or more). T may be a polyclonal antibody, in case the use is non-therapeutical.

15 It is not imperative for T to have a polypeptide structure.

The modified superantigen SA(m) is primarily a mutated superantigen but may potentially also be a chemically modified superantigen, including fragments of superantigens retaining the ability to bind to the V β chain of the T cell receptor.

20 The expression "mutated superantigen" means that the native ability of the superantigen to bind to MHC class II antigens has been modified on the genomic level by replacing, inserting or removing one or more amino acids in the native superantigen.

Superantigen fragments obtained by mutations removing parts of 25 the full amino acid sequence and fragments obtained by enzymatic or chemical cleavage of superantigens may be used equivalently in chemical conjugates of the invention.

The modified superantigen SA(m) may comprise one or more amino acid sequences that are derived from different superantigens and 30 that may have been mutated, for instance combinations of the preferred superantigens mentioned below.

The modified superantigen SA(m) as such may exhibit a decreased immunogenicity and toxicity compared to the native superantigen.

35 Other groups/substances that are capable of cross reacting with the V β -chain of the T cell receptor may potentially also be

employed equivalently with the mutated superantigen (SA(m)) as given above. Such groups/substances may be of non-polypeptide structure.

At the end of the priority year the most interesting product candidates of the invention comprised mutated forms of superantigens having multiple MHC class II binding sites and/or the ability to coordinate Zn^{2+} , for instance SEA, SED, SEE and SEH.

T as well as SA(m) may be prepared by recombinant techniques.

The bridge B may be selected as previously described (Dohlsten et al., WO 9201470), i.e. it shall preferably be hydrophilic and exhibit one or more structure(s) selected among amide, thioether, ether, disulfide etc. In case the bridge have unsubstituted unbroken hydrocarbon chains they preferably lack aromatic rings, such as phenyl. The most important bridges are those obtained by recombinant techniques, i.e. when the conjugation takes places on the genomic level. In such cases oligopeptide bridges containing hydrophilic amino acid residues, such as Gln, Ser, Gly, Glu and Arg, are preferred. Pro and His may also be included. During the priority year it has been decided that the preferred bridge is a peptide comprising three amino acid residues (GlyGlyPro).

The inventive conjugate may comprise one or more modified superantigen(s) per biospecific affinity counterpart and vice versa. This means that T in the formula above may contain one or more modified superantigens in addition to the biospecific counterpart. In analogy SA(m) may contain one or more biospecific affinity counterpart(s) T. The affinity counterpart T and SA(m) may also comprise other structures. The number of modified superantigens per affinity counterpart is preferably one or two.

The synthesis of the novel inventive conjugates may be carried out in principle according to two main routes: 1. by recombinant techniques and 2. chemical linking of T to SA(m). The methods are well recognized for the ordinary skilled worker in the field and comprise a large number of variants. It follows that the invention primarily concerns artificial conjugates, i.e. conjugates that are not found in nature.

Chemical linking of a modified superantigen to the biospecific affinity counterpart T often utilizes functional groups (e.g. primary amino groups or carboxy groups) that are present at many positions in each compound. It follows that the final product
5 will contain a mixture of conjugate molecules differing with respect to the position at which linking has taken place.

For recombinant conjugates (fusion proteins) the obtained conjugate substance will be uniform with respect to the linking position. Either the amino terminal of the modified superantigen
10 is linked to the carboxy terminal of the biospecific affinity counterpart or vice versa. For antibodies, such as intact antibodies and antigen binding fragments (Fab, Fv etc), either the light or the heavy chain may be utilized for such fusions. At present time recombinant conjugates are preferred, with
15 preference for Fab fragments and linking of the amino terminal of the modified superantigen to the first constant domain of the heavy antibody chain (CH1), without exclusion of the analogous linking to the light chain or to the VH and VL domain that also may give quite good results.

20 There are two different methods for obtaining large amounts of superantigens (including modified and fused forms) in E. coli: intracellular production or secretion. The latter method is preferred for the inventive conjugates because it offers purification of correctly folded protein from the periplasma and
25 from the culture medium. Intracellular production results in a complicated purification procedure and often needs refolding in vitro of the protein (in order for the protein to obtain the correct tertiary structure). The above does not exclude that it is possible to produce active conjugates also in other host
30 cells, e.g. eukaryotic cells, such as yeast or mammalian cells.

The production of mutated superantigens and selection of mutants having a modified ability to bind (affinity) to MHC class II antigens may be carried out according to known techniques (see e.g. Kappler et al., J. Exp. Med. 165 (1992) 387-396). See also
35 our experimental part.

The ability of the conjugate to bind to the T cell receptor V β chain, to the target structure and to cause lysis of the target cell depends on i.a. the peptide (SA(m)) that is derived from a superantigen, the biospecific affinity counterpart (T) and the structure and length of the bridge (B). A person ordinary skilled in the art is able to optimize the inventive conjugates with respect to the binding ability and the ability to cause lysis by studying the relationship between effect and structure with the aid of those models that have been disclosed in connection with previously known superantigen antibody conjugates (see the above-referred publications). See also the experimental part below.

By modified ability to bind MHC class II antigens is primarily intended that the ratio $IC_{50}(SA(wt)):IC_{50}(SA(m))$ is < 0.9 (90 %), such as < 0.5 (< 50 %) and possibly also < 0.01 (< 1 %). In the alternative the modified binding ability of the inventive conjugates can be measured as the ratio of the dissociation constants $K_d(SA(wt)):K_d(SA(m))$ with K_d measured in nM and with the same limits as for the ratio $IC_{50}(SA(wt)):IC_{50}(SA(m))$. For the determination of $IC_{50}(SA(wt))$, $IC_{50}(SA(m))$, $K_d(SA(m))$ and $K_d(SA(m))$ see the experimental part below.

It is previously known that certain superantigens may have two or more sites that bind to MHC class II antigen (Fraser et al., In: Superantigens: A pathogens view on the immune system. Eds. Huber & Palmer, Current Communications in Cell Molecular Biology 7 (1993) 7-29). For this type of superantigens the binding ability shall be modified at least one of the binding sites, e.g. as a reduction of the above-mentioned size. Possibly it may suffice with a superantigen modification that create a changed difference in affinity for two MHC class II binding sites, tentatively > 10 % and preferably by reducing the affinity of at least one site.

Superantigens bind to TCR V β chains of different subgroups with varying affinities. In the inventive fusion proteins/conjugates, the superantigen employed may have been modified so as to show an altered subgroup specificity or an altered affinity to one or more members of the subgroup. There

are strong reasons to believe that a parabolic relationship exists between the affinity for TCR V β and stimulation via TCR, i.e. a moderate affinity will give the maximal stimulation. Accordingly an appropriate affinity of a modified superantigen for TCR V β may be at hand as soon as the fusion protein/conjugate comprising the modified superantigen is able to significantly stimulate a resting T cell population representing essentially the distribution of all human V β subgroups to proliferate. The T cell population may be pooled T cells from randomly selected human individuals. By significantly is meant that the stimulation is possible to measure. The results presented in Table II (right column) in the experimental part indicate that the ability to cause SADCC of the inventive conjugates/fusion proteins often is essentially the same as for the fusion comprising the wild-type superantigen.

Main use of the conjugates/fusion proteins of the invention.

The conjugates according to the invention are primarily intended for the treatment of the same diseases as the conjugates between normal superantigens and antibodies. See the above-mentioned publications. Thus the inventive conjugates may be administered either as the main therapy or as adjuvant therapy in connection with surgery or other drugs.

The pharmaceutical composition of the invention comprises formulations that as such are known within the field but now containing our novel conjugate. Thus the compositions may be in the form of a lyophilized particulate material, a sterile or aseptically produced solution, a tablet, an ampoule etc. Vehicles such as water (preferably buffered to a physiologically pH value by for instance PBS) or other inert solid or liquid material may be present. In general terms the compositions are prepared by the conjugate being mixed with, dissolved in, bound to, or otherwise combined with one or more water-soluble or water-insoluble aqueous or non-aqueous vehicles, if necessary together with suitable additives and adjuvants. It is imperative that the vehicles and conditions shall not adversely affect the activity

of the conjugate. Water as such is comprised within the expression vehicles.

Normally the conjugates will be sold and administered in predispensed dosages, each one containing an effective amount of the conjugate that, based on the result now presented, is believed to be within the range of 10 μ g - 50 mg. The exact dosage varies from case to case and depends on the patient's weight and age, administration route, type of disease, antibody, superantigen, linkage (-B-) et.

The administration routes are those commonly known within the field, i.e. a target cell lysing effective amount or a therapeutically effective amount of a conjugate according to the invention is contacted with the target cells. For the indications specified above this mostly means parenteral administration, such as injection or infusion (subcutaneously, intravenously, intra-arterial, intramuscularly) to a mammal, such as a human being. The conjugate may be administered locally or systemically.

By "target cell lysing effective amount" is contemplated that the amount is effective in activating and directing T-lymphocytes to destroy the target cell.

At the end of the priority year it had been decided that the preferred administration route for conjugates/fusion proteins comprising unmodified superantigens is 3 hours' intravenous infusion per day combined with a fever-reducing agent (paracetamol). The administration is to be repeated during 4 days and stopped before dsecondary antibodies are raised against the fusion protein/conjugate in the patient. This dosage schedule is likely to be applicable also to the present inventive conjugates/fusion proteins.

Alternative fields of use.

The inventive conjugates can also be employed to quantitatively or qualitatively detect the structure against which the target-seeking group (T) is directed. In general these methods are well-known to people in the field. Thus, the modified superantigen may function as a marker group within immunoassays

including immunohistochemistry meaning that the marker group in turn is detected by for instance an antibody that is directed towards the peptide (SA(m)) and labelled with an enzyme, isotope, fluorophor or some other marker group known per se. Another immunoassay method is to detect in a cell population cells that on their surface express a structure capable of binding to the target-seeking group (T). This use means that a sample from the cell population is incubated with T-lymphocytes together with the present inventive conjugate as in an SADCC assay. In case the incubation leads to cell lysis this is an indication that the population contains cells that on their surface express the structure.

EXPERIMENTAL PART

15 **MANUFACTURE OF RECOMBINANT PROTEINS**

Antibodies

The experimental work in connection with the invention has primarily been done with monoclonal antibody C215 as a model substance. This antibody is directed against an antigen in the GA-733 family (see for instance EP 376,746) and references cited therein and Larsson et al., Int. J. Canc. 32 (1988) 877-82). The C215 epitope has been judged not to be sufficiently specific for cancer treatment in humans. At the priority date mab C242 (Lindholm et al., WO 9301303) was believed to be a better candidate, as judged from experiments with its fusion product with wild-type SEA.

Bacterial strains and plasmids

The E. coli strains UL635 (xyl-7, ara-14, T4^R, ΔompT) and HB101 (Boyer and Roulland-Dessoix, J. Mol. Biol. 41 (1969) 459-472) were used for the expression and cloning, respectively. The vector pKP889 was used for expression of Fab-SEA fusion proteins (derived from the murine antibody C215) and the vectors pKP943 and pKP1055 for secretion of SEA (Fig 1). The Fab-SEA expression vector pKP889 is identical to pKP865 (Dohlsten et al, Proc. Natl. Acad. Sci. USA (1994) in press) except that the spacer between

C_H1 and SEA is GlyGlyAlaAlaHisTyrGly. Expression from pKP943 yields SEA with the native amino terminus. The use of pKP1055 results in SEA having a Gly residue added at the amino terminus. In both vectors the signals from staphylococcal protein A (Uhlén et al., J. Biol. Chem. 259 (1984) 1695-1702) are used for transcription and translation and a synthetic signal peptide for secretion (L. Abrahmsén, unpublished).

In vitro mutagenesis

10 Mutations were made by polymerase chain reactions run on a Perkin Elmer Thermocycler. The reaction mixture (100 µl) contained: 1 x PCR buffer from Perkin Elmer Cetus (10 mM Tris/HCl pH 8.3, 1.5 mM MgCl₂, 0.001 % (w/v) gelatine, an additional 2 mM MgCl₂, 0.4 mM dNTPs (Perkin Elmer Cetus), 2.5 units of Ampli Taq
15 DNA polymerase (Perkin Elmer Cetus, USA) and 100 ng DNA template. Primers were added to a final concentration of 0.8 µM. The original template was a plasmid containing Staphylococcus aureus enterotoxin A gene identical to the one published by Betley et al. (J. Bacteriol. 170 (1988) 34-41), except that the first codon
20 (encoding Ser) was changed to TCC to furnish a Bam HI site at the 5' end of the gene. Later a derivative containing more unique restriction enzyme sites introduced by silent mutations was used. Mutations introduced next to a restriction site were made with one set of primers, one of these spanning the mutation and the
25 restriction site. For most mutations two set of primers had to be used and the PCR was performed in two consecutive steps. A new restriction enzyme site was introduced together with each mutation to enable facile identification. Oligonucleotides used as primers were synthesized on a Gene Assembler (Pharmacia
30 Biotech AB, Sweden). To confirm each mutation the relevant portion of the nucleotide sequence was determined on an Applied Biosystems DNA-Sequenser using their Taq DyeDeoxy Termination Cycle Sequencing Kit.

Protein production and analysis

E. coli cells harboring the different gene constructs were grown overnight at room temperature (Fab-SEA vectors) and at 24-34°C (secretion vectors, the optimum depends on the mutation).

- 5 The broth was 2 x YT (16 g/l Bacto trypton, 10 g/l Bacto yeast extract, 5 g/l NaCl) supplemented with kanamycin (50 mg/l). Fusion proteins were induced by addition of isopropyl- β -D-thiogalactoside to a final concentration of 100 μ M. (The protein A promotor used in the expression of non-fused SEA is
- 10 constitutive). The cells were pelleted at 5000 x g and the periplasmic contents were released by gently thawing the previously frozen cell pellet in 10 mM Tris-HCl (pH 7.5) on ice during agitation for 1 hour. The periplasmic extracts were clarified by centrifugation at 9500 x g for 15 minutes. The Fab-
- 15 SEA proteins were used without further purification. SEA and Gly-SEA were further purified by affinity chromatography on an anti-SEA antibody column. Polyclonal rabbit anti-SEA antibodies were previously collected from rabbits preimmunized with SEA and purified by affinity chromatography on protein G Sepharose®
- 20 (Pharmacia Biotech).

Protein Analysis

- The proteins were separated in precast polyacrylamide SDS Tris-Glycine Novex gels (gradient 4-20 % or homogenous 12 %,
- 25 Novex novel experimental technology) and either stained with Coomassie Blue or used in Western blot. Polyclonal rabbit anti-SEA antibodies (above) were used to detect SEA in Western blot analysis, followed by porcine anti-rabbit Ig antibodies, and rabbit anti-horseradish peroxidase antibodies and peroxidase.
- 30 With Fab-SEA fusion proteins peroxidase conjugated rat antibodies recognizing the kappa chain were also used (AAC 08P, Serotech LTD, England). 3,3'-diaminobenzidine (Sigma) was used for visualization of peroxidase.

- Circular dichroism (CD) spectra were collected in a J-720
- 35 spectropolarimeter (JASCO, Japan) at room temperature (22-25°C) in 10 mM phosphate buffer, pH 8.2, with 0.02 mM ZnSO₄ and 0.005 %

(v/v) Tween[®] 20. The scanning speed was 10 nm/min and each spectrum was averaged from five subsequent scans. The cell path length was 1 mm and the protein concentration 0.2 to 0.5 mg/ml. Guanidine hydrochloride (Gdn-HCl) denaturations at equilibrium were measured at 23°C by CD at 222 nm with a protein concentration of 0.3 mg/ml and a cell path length of 1 mm. These data were used to calculate the apparent fraction of unfolded protein (F_{app}). Equilibrium unfolding parameters were derived by fitting the data to a two-site folding process (Hurle et al., Biochemistry 29 (1990) 4410-4419).

BINDING AND FUNCTIONAL ASSAYS IN VITRO

Materials

Reagents: RPMI 1640 medium obtained from Gibco, Middlesex, England was used. The medium had a pH of 7.4 and contained 2 mM L-glutamine (Gibco, Middlesex, England), 0.01 M HEPES (Biological Industries, Israel), 1 mM NaHCO₃ (Biochrom AG, Germany), 0.1 mg/ml Gentamycin sulphate (Biological Industries, Israel), 1 mM Na-pyruvate (JRH Biosciences Industries, USA), 0.05 mM mercaptoethanol (Sigma Co., USA), 100 times concentrated non-essential amino acids (Flow Laboratories, Scotland) and was supplemented with 10 % fetal bovine serum (Gibco, Middlesex, England). Recombinant SEA(wt), SEA(m) and the fusion products C215Fab-SEA(wt) and C215Fab-SEA(m) were obtained as described above. Human recombinant IL-2 was from Cetus Corp., USA. Mitomycin C was from Sigma Co., USA. Na₂⁵¹CrO₄ was obtained from Merck, Germany. Phosphate buffered saline (PBS) without magnesium and calcium was received from Imperial, England.

Cells: The human colon carcinoma cell line Colo205 and the B cell lymphoma cell line Raji were obtained from American Type Cell Culture Collection (Rockville, MD, USA) (expressing HLA-DR3/w10, -DP7, -DQw1/w2). The EBV-transformed lymphoblastoid B cell line BSM was a generous gift from Dr van De Griend, Dept of Immunology, Dr Daniel den Hoed Cancer Center, Leiden, the Netherlands. The cells were repeatedly tested for mycoplasma

contamination with Gen-Probe Mycoplasma T.C. test, Gen-Probe Inc., San Diego, USA.

SEA activated T cell lines were produced by activation of mononuclear cells from peripheral blood. The blood was received
5 as buffy coats from blood donors at the University Hospital of Lund. The PBMs were stimulated at a concentration of 2×10^6 cells/ml with mitomycin C treated SEA coated BSM cells (preincubated with 100 ng/ml SEA) in medium with 10% FCS. The T cell lines were restimulated biweekly with 20 U/ml human
10 recombinant IL-2 and weekly with mitomycin C treated SEA coated BSM cells. The cell lines were cultivated for 4-12 weeks before being used in the assay.

The viability of the effector cells, as determined by trypan blue exclusion, exceeded 50 %.

15

Determination of MHC class II binding characteristics of wild-type and mutant SEA

Radioiodination procedure. Appropriate amounts of wild-type or mutant SEA were radiolabeled with 10 to 25 mCi Na^{125}I using
20 enzymobeads with the lactoperoxidase technique (NEN, Boston, MA). The reaction was stopped by quenching with sodium azide and protein-bound radioactivity was separated from free iodine by filtration through a PD-10 column (Pharmacia Biotech AB, Sweden) with R10 medium as elution buffer. Conditions were chosen to
25 obtain a stoichiometric ratio between iodine-125 and protein of $\leq 2:1$. The radiochemical purity was verified by size-exclusion chromatography on a TSK SW 3000 HPLC column. The effect of the radioiodination on the binding activity was only tested for wild-type SEA and found not to be affected (data not shown).

30 **Direct binding assay.** Raji cells, $6 \times 10^4/100 \mu\text{l}$, previously cultivated in R10 medium, were added to conical polypropylene tubes and incubated ($22^\circ\text{C}/45 \text{ min}$) in triplicate with 100 $\mu\text{l}/\text{tube}$ of serially diluted ^{125}I -labeled wild-type or mutant SEA. The cells were washed with 2 ml 1 % (w/v) bovine serum albumin
35 (BSA) in 10 mM phosphate-buffered saline (PBS), pH 7.4, centrifugated at $300 \times g$ for 5 minutes and aspirated. This

procedure was repeated twice. Finally, the cells were analyzed for cell-bound radioactivity in a gamma counter (Packard Instruments Co, Downers Grove, IL, USA). The apparent dissociation constant, K_d , and the number of binding sites, N , at saturation were calculated according to Scatchard (Ann. N.Y. Acad. Sci. 51 (1949) 660-72) after subtraction of non-specific binding (i.e. binding after incubation with R10 medium alone.

Inhibition assay (inhibition of ^{125}I -labeled wild-type SEA binding by mutant SEAs). These inhibition experiments were carried out as is described for the direct binding assay with slight modifications. Briefly, 50 μl of ^{125}I -labeled wild-type SEA was allowed to compete with an excess of unlabeled wild-type or mutant SEA (50 μl /tube) for binding to 6×10^4 /100 μl Raji cells. A tracer concentration yielding $\approx 40\%$ bound radioactivity in the direct assay was used to obtain maximal sensitivity in the inhibition assay. The displacement capacity of the competitor was expressed as the concentration yielding 50% inhibition (IC_{50}) of bound radioactivity. The binding affinity of the mutants relative to wild-type SEA was calculated using the equation:

$$\text{IC}_{50}(\text{SEA}(\text{wt})) : \text{IC}_{50}(\text{SEA}(\text{m}))$$

In order to analyze whether the mutants compete for binding to the same site on Raji cells as wild-type SEA, the binding data obtained with SEA mutants were plotted as a log-logit function and tested for parallelism with the corresponding data for wild-type SEA.

Inhibition assay (inhibition of the binding of fluorescent-labeled wild-type SEA by unlabeled wild-type SEA and SEA mutants). Raji cells (2.5×10^5) were incubated with inhibitor (wild-type or mutant SEA; 0-6000 nM) diluted in 50 μl CO_2 -independent medium (Gibco) supplemented with 10% FCS, glutamine and gentamycin at 37°C for 30 minutes. Fluorescein conjugated wild-type SEA was added to a final concentration of 30 nM and the samples were incubated for an additional half hour at 37°C . The samples were washed three times with ice cold PBS supplemented with 1% BSA (PBS-BSA) and finally kept in 0.4 ml PBS-BSA on ice until they were analyzed. From each sample 10 000 live cells were

analyzed for green fluorescence on a FACStar® (Becton Dickinson) flow cytometer and the mean fluorescence value was calculated using the LYSIS II program.

5 **SDCC and SADCC assays of SEA(wt), SEA(m) and their fusion proteins with C215Fab.**

SDCC-assays. The cytotoxicity of SEA(wt), SEA(m) and their fusions with C215Fab against MHC class II⁺ Raji cells was analyzed in a standard 4 hour ⁵¹Cr³⁺-release assay, using in vitro stimulated SEA specific T cell lines as effector cells. Briefly, ⁵¹Cr labeled Raji cells were incubated at 2.5 x 10³ cells per 0.2 ml medium (RPMI, 10 % FCS) in microtitre wells at defined effector to target cell ratio in the presence or absence (control) of the additives. Percent specific cytotoxicity was calculated as 100 x ([cpm experimental release - cpm background release]/[cpm total release - cpm background release]). The effector to target cell ratio was 30:1 for unfused SEAs and 40:1 for fusion proteins.

SADCC against of human colon cancer cells. The cytotoxicity of C215Fab-SEA(wt), C215Fab-SEA(m), SEA(wt) and SEA mutants against C215⁺ MHC class II⁻ colon carcinoma cells SW 620 was analyzed in a standard 4 hour ⁵¹Cr³⁺-release assay, using in vitro stimulated SEA specific T cell lines as effector cells. Briefly, ⁵¹Cr³⁺-labeled SW 620 cells were incubated at 2.5 x 10³ cells per 0.2 ml medium (RPMI, 10 % FCS) in microtitre wells at effector to target cell ratio 30:1 in the presence or absence (control) of the additives. Percent specific cytotoxicity was calculated as for SDCC assays.

30 **IN VIVO FUNCTIONAL EXPERIMENTS**

Tumor cells. B16-F10 melanoma cells transfected with a cDNA encoding the human tumor associated antigen C215 (B16-C215) (Dohlsten et al., Monoclonal antibody-superantigen fusion proteins: Tumor specific agents for T cell based tumor therapy; Proc. Natl. Acad. Sci. USA, In press, 1994), were grown as adherent cells to subconfluency. The culture medium consisted of

RPMI 1640 (GIBCO, Middlesex, UK) supplemented with 5×10^{-5} β -mercaptoethanol (Sigma, St Louis, MO, USA), 2 mM L-glutamine (GIBCO), 0.01 M Hepes (Biological Industries, Israel) and 10 % fetal calf serum (GIBCO). The cells were detached by a brief incubation in 0.02 % EDTA and suspended in ice cold phosphate buffered saline with 1 % syngeneic mouse serum (vehicle) to 4×10^5 cells/ml.

Animals and animal treatment. The mice were 12-19 weeks old C57B1/6 mice transgenic for a T cell receptor $V\beta 3$ chain (Dohlsten et al., Immunology 79 (1993) 520-527). One hundred thousand B16-C215 tumor cells were injected i.v. in the tail vein in 0.2 ml vehicle. On day 1, 2 and 3, the mice were given i.v. injections of C215Fab-SEA(wt) or C215Fab-SEA(D227A) in 0.2 ml vehicle at doses indicated in the figures 5a and 5b. Control mice were given only vehicle according to the same schedule. On day 21 after tumor cell injection, the mice were killed by cervical dislocation, the lungs removed, fixed in Bouin's solution and the number of lung metastases counted.

RESULTS

"Alanine scanning" of staphylococcal enterotoxin A.

Initially the structure of SEA was unknown and only speculations could be done about what side chains were surface accessible. Therefore, the majority of the mutants were chosen from alignments of homologous superantigens (Marrack and Kappler, Science 248 (1990) 705-711). Conserved (mainly polar) residues were chosen on the rational that some of these superantigens are expected to bind to HLA-DR in a conserved fashion (Chitagumpala et al., J. Immunol. 147 (1991) 3876-3881). Alanine replacements were used according to published strategies (Cunningham and Wells, Science 244 (1988) 1081-1085). During the course of this work the available information increased: i) it was shown that a Zn^{2+} ion is important for the interaction between SEA and MHC class II (HLA-DR) (Fraser et al., Proc. Natl. Acad. Sci. USA 89 (1991) 5507-5511), ii) a mutational analysis of staphylococcal enterotoxin B (SEB) was presented (Kappler et al., J. Exp. Med.

175 (1992) 387-396), and iii) the structure of SEB was presented (Swaminathan et al., Nature 359 (1992) 801-806).

Our first mutant showing a severely reduced affinity for HLA-DR, D227A, was found to co-ordinate the Zn^{2+} ion very poorly (data not shown). Assuming a common fold for SEA and SEB, the new data suggested two MHC class II binding regions; one involving the Zn^{2+} ion and one corresponding to the site defined in SEB. A second set of mutations were made on these assumptions. This second set of mutants were expressed in the form of SEA carrying a glycine added at the amino terminus. First the extension was shown to have no effects on the binding properties of wild-type SEA (next section).

Most of the mutants were expressed and secreted by E. coli in a functional form as judged by analysis of the binding of monoclonal antibodies (Table I). Very low amounts were obtained of the mutants E154A/D156A and R160A. Consequently these were excluded from the study. The mutants having an Ala substitution in residues 128, 187, 225 or 227 were not recognized by the monoclonal antibody 1E. The latter two mutants showed a reduced level of expression (more pronounced at 34°C than at 24°C) and migrated faster during SDS-PAGE, under denaturing but not reducing conditions (all other mutants migrated as wild-type SEA, data not shown). As judged by CD spectra analysis the structure of D227A could differ slightly from native SEA (figure 2), but the stability was very close to wild-type SEA (measured as resistance towards guanidine hydrochloride denaturation). The calculated $\Delta\Delta G$ between the mutant and native SEA (SEA(wt)) was - 0.16 kcal/mol and is only about 4 % of the ΔG° values (data not shown). Overall the signals in the CD analysis were low, as expected from a mostly β -sheet structure. It was recently reported that His 225 co-ordinates Zn^{2+} (unpublished data in Fraser et al (Proc. Natl. Acad. Sci. USA 89 (1991) 5507-5511). Since Asp 227 is involved in Zn^{2+} co-ordination (above) and presumably located in the same β -sheet as His 225 this suggests that these two residues constitutes the zinc-binding nucleus

found in zinc-co-ordinating proteins (Vallee and Auld, Biochemistry 29 (1990) 5647-5659).

Binding to MHC class II and T cell receptor

5 The MHC class II affinity was calculated from the amounts needed to compete with fluorescein-labeled wild-type SEA for Raji cell exposing large amounts of MHC class II. The displacement capacity of a mutant was calculated from the concentration yielding 50 % inhibition (IC₅₀) of bound fluorescence compared
10 with the concentration needed with wild-type SEA as the competitor. For wild-type SEA and for some mutants, the result from this analysis was compared with the result from an analysis where ¹²⁵I labeled wild-type SEA was used as the tracer. As may be seen in Table II, the values obtained from these two
15 inhibition analyses correlate well

For six selected mutants the binding to MHC class II was measured directly using ¹²⁵I labeled mutant SEA (Table II). With the mutant H50A the values obtained from the direct binding assay and the inhibition assays correlated well but with the mutant
20 F47A a large discrepancy was found: the direct binding indicated only 7 times weaker binding than wild-type SEA but both competition analyses demonstrated around 70 times reduced binding. The data from two of the other mutants indicated two separate binding interactions. For the mutants H225A and D227A
25 the affinity was below the detection limit also in this analysis.

We previously showed that fusion proteins composed of the Fab fragment of a carcinoma reactive antibody and SEA could be used to direct cytotoxic T cells to specifically lyse cancer cells, while the interaction between SEA and the T cell receptor (TCR)
30 was too weak to be detected by itself (Dohlsten et al., Proc. Natl. Acad. Sci. USA, in press). Thus, in contrast to analyses involving the isolated superantigen the Fab fusion context enables a functional assay for the interaction between SEA and the TCR, independent of the MHC class II binding. Consequently,
35 the efficiency of the different conjugates to direct T cells to lyse cells recognized by the Fab moiety was monitored in a

chromium release assay. This analysis confirmed that the mutations shown to affect the MHC class II binding did not affect the TCR binding (Table II).

5 **Biological effects of the mutations**

The proliferative effect was measured as the ability to stimulate peripheral lymphocytes to divide. All three mutants that competes very poorly for MHC class II induced little or no proliferation and the intermediate mutant H187A displayed some proliferative capacity, whereas the other investigated mutants were indistinguishable from the wild-type (table III). Harris et al (Infect. Immun. 61 (1993) 3175-3183) recently reported a similar severe reduction in T cell stimulatory activity for the SEA mutants F47G and L48G. Clearly a strong reduction in any of the two suggested binding regions results in a severe effect on the ability to induce proliferation. This suggests that SEA cross-links two molecules of MHC class II leading to dimerization of the TCR and that this is needed to yield a signal transduction.

In contrast the efficiency of the different mutants in directing in vitro stimulated SEA T cells to lyse MHC class II bearing target cells shows correlation with the binding affinity, rather than to the ability to compete (Table III). For example, the efficiency of F47A and D227A are only reduced 2.5 times and 300 times, respectively. Thus, here no inherent requirement for divalency too is obvious. The increase in multivalency resulting from the significantly larger number of TCRs on the surface of activated T cells might partially shield the effect of a lower avidity in the SEA/MHC class II interaction. That dimerization is not needed to direct T cell cytotoxicity has previously been demonstrated by the use of carcinoma specific bifunctional antibodies containing one anti-CD3 moiety and one anti-carcinoma moiety (Renner et al., Science 264 (1994) 833-35).

In vivo functional experiments: The results are represented in figures 6a and 6b. Treatment of mice with C215Fab-SEA(wt) and

C215Fab-SEA(D227A) were both highly effective in reducing the number of lung metastases of B16-C215 melanoma cells. The therapeutic effect was essentially identical for the two variants of the targeted superantigens. Treatment with C215Fab-SEA(wt) resulted in 70 % lethality at doses of 5 µg/injection. In contrast, no mice died when the same dose of C215Fab-SEA(D227A) were used. Taken together, SEA(D227A) is an example of a mutant with reduced toxicity and retained therapeutic efficiency when incorporated in a Fab-SEA fusion protein.

DISCUSSION

The structure of the complex between SEB and HLA-DR was recently reported (Jardetzky et al., Nature 368 (1994) 711-718). Most of the SEB residues identified to be involved in this interaction are conserved in SEA. Our data on mutant D227A indicates a weak affinity for the interaction between this site of SEA (the amino proximal site) and the MHC class II, having a K_d value higher than 8 µM. The K_d for the interaction between SEB and HLA-DR was recently reported to be 1.7 µM (Seth et al., Nature 369 (1994) 324-27). The different interactions between SEB, TCR and HLA-DR were investigated and it was shown that the complex between SEB and HLA-DR was not stably maintained in the absence of TCR. Plasmon resonance experiments indicated that this was because of a very fast off-rate. The avidity effects obtained if SEA cross-links two molecules of MHC class II followed by a subsequent dimerization of the TCR could explain how SEA may induce proliferative effects at concentrations well below the K_d . Assuming that the mutation F47A reduces the affinity of the amino proximal site below significance, the K_d of the Zn^{2+} site is around 95 nM. This hypothesis was recently strengthened by the observation that the mutants F47R, F47R/H50A and F47R/L48A/H50D show identical affinity for MHC class II as F47A (unpublished).

Based on the SEB structure (Kappler et al., J. Exp. Med. 175 (1992) 387-396) and on homology alignments (Marrack and Kappler, Science 248 (1990) 705-711), it is strongly suggested that His225 and Asp227 are located in the same β-sheet and thus the side

chains could be proximal. Thus, most likely these two residues constitute the zinc-binding nucleus found in zinc-co-ordinating proteins (Vallee and Auld, Biochemistry 29 (1990) 5647-5659). Similarly to these mutants, the mutants with a replacement at residue 128 or 187 are also recognized by all monoclonals except 1E. Fraser et al (Proc. Natl. Acad. Sci. USA 89 (1991) 5507-5511) showed that Zn^{2+} is bound to SEA and is needed for a high affinity interaction with MHC class II. The affinity for zinc was not affected by the addition of HLA-DR. Based on this observation and the high affinity for Zn^{2+} (K_d of around 1 μM) a co-ordination exclusively provided by SEA and involving 4 fold co-ordination was suggested. Our data indicates an involvement of the four residues N128, H187, H225 and D227. The function of the former two residues is not yet clear; instead of providing a ligand N128 could help in the deprotonation of D227. One argument for this is that the effect of replacing D227 is more severe than when replacing H225.

It was previously reported that there is a lack of correlation between the affinity of different superantigens for the MHC class II and the capacity to stimulate T cells to proliferate (Chintagumpala et al., J. Immunol. 147 (1991) 3876-3881). These results might partly be explained by different affinities of the superantigens towards different TCR V β -chains. Here we have observed the same lack of correlation but in contrast to separate superantigens the mutants display identical TCR affinity as shown in the Fab-SEA context (measured as SADCC). The most likely explanation for the lack of correlation is that two binding regions identified in this analysis represent two separate binding sites that yields not only a co-operative binding, but which results in the cross-linking of two molecules of MHC class II, which in turn yields dimerization of two molecules of the T cell receptor. This would imply that the affinity of both sites are important to obtain the proliferative effect. A high avidity results from the interactions within a hexameric complex involving two molecules of SEA, TCR and MHC class II. Thus the

strong affinity/avidity of SEA towards MHC class II enables SEA interaction with the TCR despite a low direct affinity.

Other biospecific affinity counterparts: A fusion protein of

5 SEA(D227A) and an IgG-binding domain of staphylococcal protein A has been produced by recombinant technology and expressed in *E. coli*. This reagent has successfully been used to target T-lymphocytes to Mot 4 and CCRF-CEM cells (obtained from ATCC) that are CD7 and CD38 positive but HLA-DP, -DQ and -DR negative. The
10 Mot 4 and CCRF-CEM cells were preincubated with anti-CD7 or anti-CD38 mouse monoclonals (Dianova, Hamburg, Germany). In order to enhance binding between the mouse monoclonals and the IgG-binding part of the fusion protein rabbit anti-mouse Ig antibody was also added.

15 In comparison with protein A-SEA(wt), protein A-SEA(D227A) had a decreased ability to bind to Daudi cells expressing MHC class II antigen.

Table I

20 Confirmation of mutant structural integrity. The binding of six monoclonal antibodies was monitored.

	Mutation	Monoclonal antibody					
		1A	2A	3A	1E	4E	EC-A1
	Wild-type	+	+	+	+	+	+
25	D11A/K14A	+	+	+	+	+	+
	D45A	+	+	+	+	+	+
	F47A	+	+	+	+	+	+
	H50A	(+)	+	(+)	+	+	+
	K55A	+	+	+	+	+	+
30	H114A	+	+	+	+	+	+
	K123A/D132G	+	+	+	+	+	+
	N128A	+	+	+	-	+	+
	K147A/K148A	+	+	+	+	-	+
	E154A/D156A	ND	ND	ND	+	ND	ND
35	R160A	ND	ND	ND	+	ND	ND
	H187A	+	+	+	-	+	+

25

E191A/N195A	+	+	+	+	+	+
D197A	+	+	+	+	+	+
H225A	.	+	+	-	+	+
D227A	+	+	+	-	+	+

- 5 Footnotes: A plus sign indicates binding, parenthesis indicate 50 to 90 % binding compared with wild-type SEA. ND means not determined.

Table II

- 10 Binding of SEA mutants to the MHC class II and the T cell receptor. The latter was monitored as the ability to direct activated cytotoxic T-cells specifically to lyse carcinoma cells using Fab-SEA fusions of the different mutants (SADCC).

Mutation	IC ₅₀ (nM) SEA-FITC ¹	IC ₅₀ (nM) 125I-SEA ¹	K _d (nM) 125I labeled ¹	SADCC(% of wild-type ¹
15 wild-type	50	38	13	100 ²
Gly-SEA	50	ND	ND	100 ²
D11A/K14A	50	ND	ND	ND
D45A	53	ND	ND	ND
20 F47A	3150	2943	95	100
H50A	150	132	32	100
K55A	44	ND	ND	ND
H114A	48	ND	ND	ND
K123A/D132G	188	75	12/237	100
25 N128A	1150	ND	2.9/76	100
K147A/K148A	58	ND	ND	ND
H187A	1030	602	97	100
E191A/N195A	51	ND	ND	ND
D197A	78	ND	ND	ND
30 H225A	>9000	9600	ND	ND
D227A	>9000	>10000	>8000	100

Footnotes: 1) ND means not determined. 2) In the Fab-SEA context the spacer between C_H1 and SEA ends with a Gly.

Table III

Biological effects of the mutations. The ability to stimulate resting T cells to proliferate and the ability to direct cytotoxic cells to lyse MHC class II exposing target cells were monitored (SDCC = Superantigen Dependent mediated Cellular Cytotoxicity).

	Mutation	Proliferation	SDCC
		%	EC ₅₀ (relative)
10	wild-type	100	1
	Gly-SEA	ND	1
	D11A/K14A	ND	0.8
	D45A	50	1.3
	F47A	<0.2	2.5
15	H50A	20	1.4
	K55A	100	1.3
	H114A	ND	1
	K123A/D132G	40	2.1
	N128A	40	1.2
20	K147A/K148A	ND	0.7
	E154A/D156A	ND	ND
	R160A	ND	ND
	H187A	15	4
	E191A/N195A	100	1.1
25	D197A	ND	1.3
	H225A	<0.2	3x10 ²
	D227A	<0.01	3x10 ²

Footnotes: ND means not determined.

30 LEGENDS TO THE FIGURES

General: The mutant SEA(D227A) (=SEA(m9) or mutant m9) was at the priority date the most promising SEA variant. We have therefore selected to present in vitro and in vivo results with this variant (Figures 3-6).

Figure 1.

Schematic outline of the plasmids used to express SEA and C215Fab-SEA. The coding regions and the two transcription terminators following the product genes are indicated by boxes.

- 5 The gene encoding the kanamycin resistance protein is labeled Km. lacI is the lac repressor gene. V_H and C_H1 indicates the gene encoding the F_D fragment of the heavy chain of the murine antibody C215. Likewise V_K and C_K indicates the gene encoding the kappa chain. Rop is the gene encoding the replication control
- 10 protein from pBR322. The promoters directing transcription of product genes are shown as arrows, in pKP889 the trc promotor and in the other two vectors the promotor from staphylococcal protein A (spa). The region containing the origin of replication is indicated by ori. The only difference between SEA encoded by
- 15 pKP943 and pKP1055 is a glycine residue added at the N-terminus of the latter. The SEA gene contained in the latter vector also contains more unique restriction enzyme sites, introduced by silent mutations.

Figure 2

- 20 Circular dichroism spectra for wild-type SEA and for the mutants F47A and D227A, representing the most severely reduced mutations in each MHC class II binding region. The solid line is the curve for wild-type SEA. The curves for the mutants are dotted or center, F47A respectively D227A.

- 25 **Figure 3** shows the concentration dependency of superantigen dependent mediated cellular cytotoxicity (SDCC) for SEA(wt) and SEA(D227A).

- Figure 4** shows the concentration dependency of superantigen dependent cell mediated cytotoxicity (SDCC) for C215Fab-SEA(wt) and C215Fab-SEA(D227A).
- 30

Figure 5 shows the concentration dependency of superantigen mAb dependent cell mediated cytotoxicity (SADCC) for C215Fab-SEA(wt) and C215Fab-SEA(D227A) compared to free SEA(wt).

- Figure 6a** compares the therapeutic effects obtained in C57B1/6 mice carrying lung metastases of B16-C215 melanoma cells by
- 35 treatment with C215Fab-SEA(wt) and C215Fab-SEA(D227A).

Figure 6b shows toxicity of C215-SEA(wt) and C215-SEA(D227A) for the treatments represented in figure 6a.

P A T E N T C L A I M S

1. Conjugate comprising

5 a. a biospecific affinity counterpart (target-seeking group)
 that is capable of binding to a predetermined structure,
 and

 b. a peptide that

 i. contains an amino acid sequence that is derived from
 a superantigen,

10 ii. has the ability to bind to a V β chain of a T cell
 receptor, and

 iii. has a modified ability to bind to MHC class II
 antigens compared to the superantigen from which the
 peptide is derived,

15 which parts are covalently linked together.

2. The conjugate according to claim 1, **characterized** in that

5 a. the biospecific affinity counterpart is directed towards
 a cell surface structure, and that

20 b. the conjugate has the ability to activate T-lymphocytes
 to lyse cells that exhibit the cell surface structure on
 their surface.

3. The conjugate according to any one of claims 1-2,

25 **characterized** in that the biospecific affinity counterpart
 is an antibody or an antigen binding fragment of an
 antibody.

4. The conjugate according to any one of claims 1-3,

30 **characterized** in that it is a fusion protein.

5. The conjugate according to any one of claims 1-4,

characterized in that the peptide is a mutated superantigen.

35 6. The conjugate according to any one of claims 1-5,

characterized in that the peptide is derived from a superantigen and that its ability to bind to MHC class II antigens is altered with at least 10 %.

- 5 7. The conjugate according to any one of claims 1-6, **characterized** in that the superantigen is staphylococcal enterotoxin A, B, C₁, C₂, D, or E.
- 10 8. The conjugate according to claim 7, **characterized** in that the superantigen in addition may be derived from staphylococcal enterotoxin H
- 15 9. The conjugate according to any one of claims 1-8, **characterized** in that the structure against which the biospecific affinity counterpart is directed is a structure that is expressed on the cell surface during a disease, for instance a cancer, a viral infection, an autoimmune disease or a parasitic infestation.
- 20 10. A method for the lysis of mammalian cells, **characterized** in that the cells are contacted with T-lymphocytes and a conjugate according to any one of claims 2-9 in which the biospecific affinity counterpart is directed against a surface structure on the cells that are to be lysed, said incubation being performed under conditions allowing for lyse of said cells.
- 25 11. A method for selective lysis of cells (I) that are present together with other cells (II) and that express a structure that is preferentially occurring on those cells (I) that are to be lysed, **characterized** in that the cells (I together with II) simultaneously are contacted with a conjugate according to any one of claims 2-9 in which the biospecific affinity counterpart is directed towards a surface structure on the cells (I) that are to be lysed, said contact being performed under conditions permitting lysis.
- 30 35

- 10

PTO/SB/01 (4-96)

Approved for use through 9/30/98 OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Type a plus sign (+) inside this box → +

67366 DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION <input type="checkbox"/> Declaration Submitted with Initial Filing OR <input checked="" type="checkbox"/> Declaration Submitted after Initial Filing	Attorney Docket Number	A96335DS
	First Named Inventor	Abrahmsen
	COMPLETE IF KNOWN	
	Application Number	08/765,695
	Filing Date	
	Group Art Unit	
	Examiner Name	

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

A CONJUGATE BETWEEN A MODIFIED SUPERANTIGEN AND A TARGET-SEEKING COMPOUND AND THE USE OF THE CONJUGATE

(Title of the invention)

The specification of which

☐ is attached hereto

OR

☒ was filed on (MM/DD/YYYY)

06/07/1995

as United States Application Number or PCT International

Application Number

PCT/SE95/00681

and was amended on (MM/DD/YYYY)

(if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code §119 (a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365 (a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
9402430-4	Sweden	07/11/1994	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority sheet attached hereto.

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority sheet attached hereto.


Under the Paperwork Reduction Act of 1985, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner of Patents and Trademarks, Washington, DC 20231.

Type a plus sign (+) inside this box →


DECLARATION				Page 2	
<p>I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or §365(c) of any PCT International application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.</p>					
U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)		
<input type="checkbox"/> Additional U.S. or PCT International application numbers are listed on a supplemental priority sheet attached hereto.					
<p>As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:</p>					
Name	Registration Number	Name	Registration Number		
David L. Fox	240,612	Seth M. Nehrbass	31,281		
Paul E. Krieger	25,886	Jan K. Simpson	33,283		
Charles C. Garvey, Jr.	27,889	7			
Charles M. Cox	29,057				
Gregory C. Smith	29,441				
<input type="checkbox"/> Additional registered practitioner(s) named on a supplemental sheet attached hereto.					
Direct all correspondence to:					
Name	Paul E. Krieger				
Address	1177 West South Loop, 10th Floor				
Address					
City	Houston	State	TX	ZIP	77027
Country	US	Telephone	713-850-0909	Fax	713-850-0165
<p>I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.</p>					
Name of Sole or First Inventor:		<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Lars 1-00	Middle Initial		Family Name	Abrahmsen
				Suffix e.g. Jr.	
Inventor's Signature	Lars A.			Date	July 14, 1997
Residence: City	Bromma SE	State		Country	Sweden
Post Office Address	Lillangsgatan 28, S-168 58				
Post Office Address					
City	Bromma	State		Zip	
				Country	Sweden
<input checked="" type="checkbox"/> Additional inventors are being named on supplemental sheet(s) attached hereto					

Type a plus sign (+) inside this box → ☐

DECLARATION					ADDITIONAL INVENTOR(S) Supplemental Sheet	
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name	Per	Middle Initial		Family Name	Bjork	Suffix e.g. Jr.
Inventor's Signature					Date	
Residence: City	Helsingborg	State		Country	Sweden	Citizenship SE
Post Office Address	Sigridsgatan 31, S-256 56					
Post Office Address						
City	Helsingborg	State		Zip		Country Sweden
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name	Mikael	Middle Initial		Family Name	Dohlsten	Suffix e.g. Jr.
Inventor's Signature					Date	
Residence: City	Lund	State		Country	Sweden	Citizenship SE
Post Office Address	Gerdagatan 12.B, S-223 62					
Post Office Address						
City	Lund	State		Zip		Country Sweden
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name	Terie	Middle Initial		Family Name	Kalland	Suffix e.g. Jr.
Inventor's Signature					Date	080797
Residence: City	Arese ITV	State		Country	Italy	Citizenship NO
Post Office Address	Viale 26/3, 20020					
Post Office Address						
City	Arese	State		Zip		Country Italy
Name of Additional Joint Inventor, if any:					<input type="checkbox"/> A petition has been filed for this unsigned inventor	
Given Name		Middle Initial		Family Name		Suffix e.g. Jr.
Inventor's Signature					Date	
Residence: City		State		Country		Citizenship
Post Office Address						
Post Office Address						
City		State		Zip		Country

☐ Additional inventors are being named on supplemental sheet(s) attached hereto

Type a plus sign (+) inside this box → ☐

DECLARATION				ADDITIONAL INVENTOR(S) Supplemental Sheet			
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Per	Middle Initial		Family Name	Bjork	Suffix e.g. Jr.	
Inventor's Signature					Date		
Residence: City	Helsingborg	State		Country	Sweden	Citizenship	SE
Post Office Address	Sigridsgatan 31, S-256 56						
Post Office Address							
City	Helsingborg	State		Zip		Country	Sweden
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Mikael	Middle Initial		Family Name	Dohlsten	Suffix e.g. Jr.	
Inventor's Signature					Date	15 July 1997	
Residence: City	Lund	State	SEX	Country	Sweden	Citizenship	SE
Post Office Address	Gerdagatan 12. B, S-223 62						
Post Office Address							
City	Lund	State		Zip		Country	Sweden
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Terje	Middle Initial		Family Name	Kalland	Suffix e.g. Jr.	
Inventor's Signature					Date		
Residence: City	Arese. ITX	State		Country	Italy	Citizenship	NO
Post Office Address	Viale 26/3, 20020						
Post Office Address							
City	Arese	State		Zip		Country	Italy
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name		Middle Initial		Family Name		Suffix e.g. Jr.	
Inventor's Signature					Date		
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	

☐ Additional inventors are being named on supplemental sheet(s) attached hereto

200

300

6696275050200

100

Type a plus sign (+) inside this box → ☒

DECLARATION				ADDITIONAL INVENTOR(S) Supplemental Sheet			
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Per	Middle Initial	Family Name	Bjork	Suffix e.g. Jr.		
Inventor's Signature	<i>Per Dan</i>			Date	July 14, 1997		
Residence: City	Helsingborg	State	Country	Sweden	Citizenship	SE	
Post Office Address	Sigridsgatan 31, S-256 56						
Post Office Address							
City	Helsingborg	State	Zip	Country	Sweden		
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Mikael	Middle Initial	Family Name	Dohlsten	Suffix e.g. Jr.		
Inventor's Signature				Date			
Residence: City	Lund	State	Country	Sweden	Citizenship	SE	
Post Office Address	Gerdagatan 12 B, S-223 62						
Post Office Address							
City	Lund	State	Zip	Country	Sweden		
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name	Terje	Middle Initial	Family Name	Kalland	Suffix e.g. Jr.		
Inventor's Signature				Date			
Residence: City	Arese	State	Country	Italy	Citizenship	NO	
Post Office Address	Viale 26/3, 20020						
Post Office Address							
City	Arese	State	Zip	Country	Italy		
Name of Additional Joint Inventor, if any:				<input type="checkbox"/> A petition has been filed for this unsigned inventor			
Given Name		Middle Initial	Family Name		Suffix e.g. Jr.		
Inventor's Signature				Date			
Residence: City		State	Country		Citizenship		
Post Office Address							
Post Office Address							
City		State	Zip	Country			

☐ Additional inventors are being named on supplemental sheet(s) attached hereto

0092055560

300

400





FIGURE 1

FIGURE 2






FIGURE 3

4/5




FIGURE 4




FIGURE 5

Therapeutic effect




FIGURE 6A

Toxicity




FIGURE 6B